

## Clinical Trials (recruiting subjects)

### ACT-1 (younger patients)

A randomized phase III study to evaluate the efficacy of chemoimmunotherapy with the monoclonal antibody Campath-1H (Alemtuzumab) given in combination with 2-weekly CHOP versus 2-weekly CHOP alone and consolidated by autologous stem cell transplant, in young patients with previously untreated systemic peripheral T-cell lymphomas

### ACT-2 (elderly patients)

A randomised phase III trial to evaluate the efficacy of chemoimmunotherapy with the monoclonal antibody Campath-1H (Alemtuzumab) given in combination with 2-weekly CHOP (Cyclophosphamide, Doxorubicin, Vincristine and Prednisone) versus 2-weekly CHOP alone in elderly patients with previously untreated systemic T- cell Lymphoma

### AIHA

Rituximab added to Prednisolone in patients with warm antibody mediated autoimmune haemolytic anaemia.

### AML17

Trial in acute myeloid leukaemia or high risk myelodysplastic syndrome 17. Protocol for patients aged under 60.

### CHEPRE / CHEPRE Nodal vs. Extranodal

A prospective validation study of chemosensitivity prediction – based on individual DNA chip analysis in patients with malignant haematological diseases requiring treatment

### CHEPRETRO

A retrospective validation of “new gene test for chemotherapy resistance” in stored tissue from patients with malignant lymphoma

### CML pregnancy registry

A global Gleevec/Glivec & Tasigna pregnancy exposure registry. A voluntary post-authorization safety study (PASS)

#### DALIAH

A Danish study of low-dose Interferon Alpha versus Hydroxyurea in the treatment of Ph negative chronic myeloproliferative neoplasms: a national randomized, prospective quality assurance study mainly focusing on efficacy, toxicity, and quality of life

#### EMN-02 / HOVON 95 MM / NMSG 19/10

A randomized phase III study to compare Bortezomib, Melphalan, Prednisone (VMP) high dose Melphalan followed by Bortezomib, Lenalidomide, Dexamethasone (VRD) consolidation and Lenalidomide maintenance in patients with newly diagnosed multiple myeloma

#### EUMDS

A prospective, multicenter European registry for newly diagnosed patients with myelodysplastic syndromes of IPSS low and intermediate-1 subtypes

#### ITP

A randomized phase III study of the efficacy of high-dose Dexamethasone versus high-dose Dexamethazone in combination with Rituximab (MabThera) in patients with newly diagnosed idiopathic thrombocytopenic purpura (ITP)

#### MSCNET MM

Cancer stem cells in multiple myeloma: a prospective registry of tissue samples

#### NIMBUS (core) Pomalidomide

A phase III, multicenter, randomized, open-label study to compare the efficacy and safety of Pomalidomide in combination with low-dose Dexamethasone versus high-dose Dexamethasone in subjects with refractory or relapsed and refractory multiple myeloma

#### NIMBUS (companion) Pomalidomide

Open-label, multi-center, single-arm study for the safety and efficacy of pomalidomide (CC-4047) monotherapy for subjects with refractory or relapsed and refractory multiple myeloma: a companion study for clinical trial CC-4047-MM-003

## MUCOSITIS

Analysis of cellular and humoral components of significance to Mucositis' pathogenesis and healing by assessment of mucosa biopsies and peripheral blood in patients treated with high dose chemotherapy

22.06.2012 /Lise Tordrup Elkjær